

Modes of transmission



Close contact (more than 15 minutes and within 2 meters) with an infected person



An infected person coughs or sneezes near you

Symptoms

- » Fever
- » Cough
- » Tiredness or fatigue
- » Loss of taste or smell
- » Shortness of breath or difficulty breathing
- » Muscle aches, sore throat, headache and/or chest pain
- » Shivering

Line of action



medical care



Stay at



Wear a face mask



Cover your nose and mouth when sneezing or coughing



Keep a distance of 2 meters from others



Wash your hands frequently



Always wash your hands before touching your face

Modes of transmission. symptoms and lines of action in patients with

sickle cell disease, hepatitis C, tuberculosis, HIV and COVID-19
























Modes of transmission



Genetic and hereditary

Symptoms

- » Anaemia
- » Pain crises / episodes
- » Swollen fingers and toes
- » Fever and bone pain
- » Vomiting or abdominal pain

Line of action



medical care



Eat a healthy diet



Rest



Drink plenty of liquids



Avoid consuming harmful substances


Avoid exposure to low oxygen levels

extreme heat

Hepatitis C

Exercise

regularly



Symptoms

- » Fever
- » Loss of appetite
- » Fatigue
- » Nausea
- » Dark urine
- » Vomiting
- »Jaundice
- » Pale stools
- » Abdominal and/or joint pain

Modes of transsmision



Unprotected sexual relations



Objects contaminated with infected blood



Blood transfusions (or blood derivatives) and organ transplants received before 1992

Line of action




Seek medical care



Avoid sharing objects that could be contaminated

Tuberculosis



Modes of transmission



Coughing, speaking or sneezing

Symptoms

- » Fatigue
- ≫ Fever
- » Night sweats
- » Shivering
- » Loss of appetite
- » Cough, with or without blood, that lasts three or more weeks
- » Chest pain when breathing or coughing
- » Involuntary weight loss

Line of action



medical care



Stay at home



Wear a face mask



Cover your nose and mouth when sneezing or coughing



Ventilate enclosed spaces



Modes of transmission

- Unprotected sexual relations
- Contaminated objects (needles and syringes)
- Blood transfusions
- Pregnancy, childbirth or breastfeeding

Symptoms

- » Fever
- » Headache, muscle and/or joint pain
- » Skin rash

- » Sore throat and painful mouth ulcers
- » Swollen lymph nodes, primarily in the neck

⊕ Asymptomatic with high risk behaviors

Line of action



medical care

___1 PLEGAR



Always use a condom





Get tested for sexually transmitted diseases



Do not inject drugs









